

Your Coronavirus Test Is Positive. Maybe It Shouldn't Be.

The usual diagnostic tests may simply be too sensitive and too slow to contain the spread of the virus.

By [Apoorva Mandavilli](#) Aug. 29, 2020



Tests authorized by the F.D.A. provide only a yes-no answer to infection, and will identify as positive patients with low amounts of virus in their bodies. Johnny Milano for The New York Times

Some of the nation's leading public health experts are raising a new concern in the endless debate over coronavirus testing in the United States: The standard tests are diagnosing huge numbers of people who may be carrying relatively insignificant amounts of the virus.

Most of these people are not likely to be contagious, and identifying them may contribute to bottlenecks that prevent those who are contagious from being found in time. But researchers say the solution is not to test less, or to skip testing people without symptoms, as recently [suggested by the Centers for Disease Control and Prevention](#).

Instead, new data underscore [the need for more widespread use of rapid tests](#), even if they are less sensitive.

"The decision not to test asymptomatic people is just really backward," said Dr. Michael Mina, an epidemiologist at the Harvard T.H. Chan School of Public Health, referring to the C.D.C. recommendation.

"In fact, we should be ramping up testing of all different people," he said, "but we have to do it through whole different mechanisms."

In what may be a step in this direction, the Trump administration announced on Thursday that it would purchase 150 million rapid tests.

The most widely used diagnostic test for the new coronavirus, called a PCR test, provides a simple yes-no answer to the question of whether a patient is infected.

But similar PCR tests for other viruses do offer some sense of how contagious an infected patient may be: The results may include a rough estimate of the amount of virus in the patient's body.

"We've been using one type of data for everything, and that is just plus or minus — that's all," Dr. Mina said. "We're using that for clinical diagnostics, for public health, for policy decision-making."

But yes-no isn't good enough, he added. It's the amount of virus that should dictate the infected patient's next steps. "It's really irresponsible, I think, to forgo the recognition that this is a quantitative issue," Dr. Mina said.

The PCR test amplifies genetic matter from the virus in cycles; the fewer cycles required, the greater the amount of virus, or viral load, in the sample. The greater the viral load, the more likely the patient is to be contagious.

This number of amplification cycles needed to find the virus, called the cycle threshold, is never included in the results sent to doctors and coronavirus patients, although it could tell them how infectious the patients are.

In three sets of testing data that include cycle thresholds, compiled by officials in Massachusetts, New York and Nevada, up to 90 percent of people testing positive carried barely any virus, a review by The Times found.

On Thursday, the United States recorded 45,604 new coronavirus cases, according to a database maintained by The Times. If the rates of contagiousness in Massachusetts and New York were to apply nationwide, then perhaps only 4,500 of those people may actually need to isolate and submit to contact tracing.

One solution would be to adjust the cycle threshold used to decide that a patient is infected. Most tests set the limit at 40, a few at 37. This means that you are positive for the coronavirus if the test process required up to 40 cycles, or 37, to detect the virus.

Tests with thresholds so high may detect not just live virus but also genetic fragments, leftovers from infection that pose no particular risk — akin to finding a hair in a room long after a person has left, Dr. Mina said.

Any test with a cycle threshold above 35 is too sensitive, agreed Juliet Morrison, a virologist at the University of California, Riverside. "I'm shocked that people would think that 40 could represent a positive," she said.

Coronavirus Schools Briefing: The pandemic is upending education. Get the latest news and tips as students go back to school.

A more reasonable cutoff would be 30 to 35, she added. Dr. Mina said he would set the figure at 30, or even less. Those changes would mean the amount of genetic material in a patient's sample would have to be 100-fold to 1,000-fold that of the current standard for the test to return a positive result.

"It's just kind of mind-blowing to me that people are not recording the C.T. values from all these tests, that they're just returning a positive or a negative," one virologist said. Erin Schaff/The New York Times

The Food and Drug Administration said in an emailed statement that it does not specify the cycle threshold ranges used to determine who is positive, and that "[commercial manufacturers](#) and [laboratories](#) set their own."

The Centers for Disease Control and Prevention said it is examining the use of cycle threshold measures "for policy decisions." The agency said

it would need to collaborate with the F.D.A. and with device manufacturers to ensure the measures “can be used properly and with assurance that we know what they mean.”

The C.D.C.'s own calculations suggest that it is extremely difficult to detect any live virus in a sample [above a threshold of 33 cycles](#). Officials at some state labs said the C.D.C. had not asked them to note threshold values or to share them with contact-tracing organizations.

For example, North Carolina's state lab uses the Thermo Fisher coronavirus test, which automatically classifies results based on a cutoff of 37 cycles. A spokeswoman for the lab said testers did not have access to the precise numbers.

This amounts to an enormous missed opportunity to learn more about the disease, some experts said.

“It's just kind of mind-blowing to me that people are not recording the C.T. values from all these tests — that they're just returning a positive or a negative,” said Angela Rasmussen, a virologist at Columbia University in New York.

“It would be useful information to know if somebody's positive, whether they have a high viral load or a low viral load,” she added.

[The Coronavirus Outbreak >](#)

Frequently Asked Questions

Updated August 27, 2020

What should I consider when choosing a mask?

There are a few basic things to consider. Does it have at least two layers? Good. If you hold it up to the light, can you see through it? Bad. Can you

blow a candle out through your mask? Bad. Do you feel mostly OK wearing it for hours at a time? Good. The most important thing, after finding a mask that fits well without gapping, is to find a mask that you will wear. Spend some time picking out your mask, and find something that works with your personal style. You should be wearing it whenever you're out in public for the foreseeable future. Read more: [What's the Best Material for a Mask?](#)

What are the symptoms of coronavirus?

In the beginning, the coronavirus [seemed like it was primarily a respiratory illness](#) — many patients had fever and chills, were weak and tired, and coughed a lot, though some people don't show many symptoms at all. Those who seemed sickest had pneumonia or acute respiratory distress syndrome and received supplemental oxygen. By now, doctors have identified many more symptoms and syndromes. In April, [the C.D.C. added to the list of early signs](#) sore throat, fever, chills and muscle aches. Gastrointestinal upset, such as diarrhea and nausea, has also been observed. Another telltale sign of infection may be a sudden, profound diminution of one's [sense of smell and taste](#). Teenagers and young adults in some cases have developed painful red and purple lesions on their fingers and toes — nicknamed "Covid toe" — but few other serious symptoms.

Why does standing six feet away from others help?

The coronavirus spreads primarily through droplets from your mouth and nose, especially when you cough or sneeze. The C.D.C., one of the organizations using that measure, [bases its recommendation of six feet](#) on the idea that most large droplets that people expel when they cough or sneeze will fall to the ground within six feet. But six feet has never been a magic number that guarantees complete protection. Sneezes, for instance, can launch droplets a lot farther than six feet, [according to a recent study](#). It's a rule of thumb: You should be safest

standing six feet apart outside, especially when it's windy. But keep a mask on at all times, even when you think you're far enough apart.

I have antibodies. Am I now immune?

As of right now, [that seems likely, for at least several months](#). There have been frightening accounts of people suffering what seems to be a second bout of Covid-19. But experts say these patients may have a drawn-out course of infection, with the virus taking a slow toll weeks to months after initial exposure. People infected with the coronavirus typically [produce](#) immune molecules called antibodies, which are [protective proteins made in response to an infection. These antibodies may](#) last in the body [only two to three months](#), which may seem worrisome, but that's perfectly normal after an acute infection subsides, said Dr. Michael Mina, an immunologist at Harvard University. It may be possible to get the coronavirus again, but it's highly unlikely that it would be possible in a short window of time from initial infection or make people sicker the second time.

I'm a small-business owner. Can I get relief?

The [stimulus bills enacted in March](#) offer help for the millions of American small businesses. Those eligible for aid are businesses and nonprofit organizations with fewer than 500 workers, including sole proprietorships, independent contractors and freelancers. Some larger companies in some industries are also eligible. The help being offered, which is being managed by the Small Business Administration, includes the Paycheck Protection Program and the Economic Injury Disaster Loan program. But lots of folks have [not yet seen payouts](#). Even those who have received help are confused: The rules are draconian, and some are stuck sitting on [money they don't know how to use](#). Many small-business owners are getting less than they expected or [not hearing anything at all](#).

What are my rights if I am worried about going back to work?

Employers have to provide [a safe workplace](#) with policies that protect everyone equally. [And if one of your co-workers tests positive for the coronavirus, the C.D.C.](#) has said that [employers should tell their employees](#) -- without giving you the sick employee's name -- that they may have been exposed to the virus.

Officials at the Wadsworth Center, New York's state lab, have access to C.T. values from tests they have processed, and analyzed their numbers at The Times's request. In July, the lab identified 794 positive tests, based on a threshold of 40 cycles.

With a cutoff of 35, about half of those tests would no longer qualify as positive. About 70 percent would no longer be judged positive if the cycles were limited to 30.

In Massachusetts, from 85 to 90 percent of people who tested positive in July with a cycle threshold of 40 would have been deemed negative if the threshold were 30 cycles, Dr. Mina said. "I would say that none of those people should be contact-traced, not one," he said.

Other experts informed of these numbers were stunned.

"I'm really shocked that it could be that high — the proportion of people with high C.T. value results," said Dr. Ashish Jha, director of the Harvard Global Health Institute. "Boy, does it really change the way we need to be thinking about testing."

Dr. Jha said he had thought of the PCR test as a problem because it cannot scale to the volume, frequency or speed of tests needed. "But what I am realizing is that a really substantial part of the problem is that we're not even testing the people who we need to be testing," he said.

The number of people with positive results who aren't infectious is particularly concerning, said Scott Becker, executive director of the

Association of Public Health Laboratories. "That worries me a lot, just because it's so high," he said, adding that the organization intended to meet with Dr. Mina to discuss the issue.

The F.D.A. noted that people may have a low viral load when they are newly infected. A test with less sensitivity would miss these infections.

But that problem is easily solved, Dr. Mina said: "Test them again, six hours later or 15 hours later or whatever," he said. A rapid test would find these patients quickly, even if it were less sensitive, because their viral loads would quickly rise.

PCR tests still have a role, he and other experts said. For example, their sensitivity is an asset when identifying newly infected people to enroll in clinical trials of drugs.

But with 20 percent or more of people testing positive for the virus in some parts of the country, Dr. Mina and other researchers are questioning the use of PCR tests as a frontline diagnostic tool.

People infected with the virus are most infectious from a day or two before symptoms appear till about five days after. But at the current testing rates, "you're not going to be doing it frequently enough to have any chance of really capturing somebody in that window," Dr. Mina added.

Highly sensitive PCR tests seemed like the best option for tracking the coronavirus at the start of the pandemic. But for the outbreaks raging now, he said, what's needed are coronavirus tests that are fast, cheap and abundant enough to frequently test everyone who needs it — even if the tests are less sensitive.

"It might not catch every last one of the transmitting people, but it sure will catch the most transmissible people, including the superspreaders,"

Dr. Mina said. "That alone would drive epidemics practically to zero."

We use cookies and similar methods to recognize visitors and remember their preferences. We also use them to measure ad campaign effectiveness, target ads and analyze site traffic. To learn more about these methods, including how to disable them, [view our Cookie Policy](#). Starting on July 20, 2020 we will show you ads we think are relevant to your interests, based on the kinds of content you access in our Services. You can [object](#). For more info, see our [privacy policy](#). By tapping 'accept,' you consent to the use of these methods by us and third parties. You can always change your tracker preferences by visiting our Cookie Policy.

[Manage Trackers](#)

Your preference has been stored for this browser and device. If you clear your cookies, your preference will be forgotten.