

BOMBSHELL: More Death and Fever from Pfizer Vaccine than Covid

William J. Banks | Paris | 9 December 2020



BREAKING: We have just seen the latest FDA report on the Phase 3 trial of Pfizer's covid vaccine (BNT162b2).

The document just released is entitled "Vaccines and Related Biological Products Advisory Committee December 10, 2020 Meeting Briefing Document" and can be downloaded [here \(.pdf\)](#). The FDA briefing document on the Pfizer-BioNTech COVID-19 vaccine contains some stunning information, including the revelation that **most people are hundreds of times more likely to have an adverse reaction from their vaccine than they are to die from covid.**

In "**Table 3. Disposition of All Randomized Participants, Phase 2/3 Safety Population**" on page 19, we find that 6 people died during the study. Pfizer tells us that 2 people died after having their Covid Vaccine and another 4 died apparently from a different vaccine, the MenACWY vaccine (which is designed to protect against meningococcal bacteria). Pfizer calls the MenACWY vaccine a 'placebo', and then use this obvious falsehood to hide the number of adverse reactions and deaths that occur.

Pfizer-BioNTech COVID-19 Vaccine
VRBPAC Briefing Document

	BNT162b2 N=18904 n (%)	Placebo N=18892 n (%)	Total N=37796 n (%)
Randomized	18904 (100.0)	18892 (100.0)	37796 (100.0)
Vaccinated			
Completed 1 dose	18858 (99.8)	18849 (99.8)	37707 (99.8)
Completed 2 doses	18555 (98.2)	18533 (98.1)	37088 (98.1)
Withdrawn from Study	180 (1.0)	259 (1.4)	439 (1.2)
Reason for Withdrawal			
Adverse Event	8 (0.0)	5 (0.0)	13 (0.0)
Death	2 (0.0)	4 (0.0)	6 (0.0)
Withdrawal by Subject	84 (0.4)	157 (0.8)	241 (0.6)
Lost to Follow-up	80 (0.4)	86 (0.5)	166 (0.4)
No longer meets eligibility criteria	1 (0.0)	2 (0.0)	3 (0.0)
Refused further study procedures	0	1 (0.0)	1 (0.0)

Source: EUA 27036, amendment 3, Table 2, c4591001-safety-tables-cos-reacto.pdf, page 43.
Note: One participant was randomized but did not sign informed consent and therefore not included in any analysis population.
Note: 120 HIV-positive participants included in this table. HIV population analyses were summarized separately from analyses based on the phase 2/3 safety population, but included in the all-enrolled population analyses presented in this briefing document.
%n/N, n = number of subjects with the specified characteristic. N = number of participants ≥16 years of age enrolled by October 9, 2020, including 120 HIV-positive participants, and received at least 1 dose of study vaccine or placebo. N is the denominator used for the percentage calculations.
Data analysis cutoff date: November 14, 2020

Obviously a **MenACWY vaccine is not a placebo**. The fact that 4 of their participants died from this vaccine and 2 from their covid vaccine only adds more weight to the argument that many vaccines are dangerous and potentially life threatening. However Pfizer skews this data by saying that 4 people died 'from a placebo'.

Let's just remind ourselves what the definition of the world 'placebo' is.

The Oxford English Dictionary has :

- Placebo – a **substance that has no physical effects**, given to patients who do not need medicine but think that they do, or used when testing new drugs"

And the Merriam-Webster Dictionary has:

- a usually **pharmacologically inert preparation** prescribed more for the mental relief of the patient than for its actual effect on a disorder
- an **inert or innocuous substance** used especially in controlled experiments testing the efficacy of another substance (such as a drug)

So for anyone who understands the English language, a MenACWY vaccine is clearly not a placebo. It would seem that Pfizer have redefined the word 'placebo' to mean 'MenACWY vaccine'. Should this type of deception be tolerated by the public? We shall leave that for you to ponder.

On Page 41 of the report, we can see a description of the Serious Adverse Events including 6 deaths and 12 cases of appendicitis. **In total 0.6% of participants reported a Serious Adverse Events from the Covid Vaccine – 100 TIMES HIGHER THAN THE PROBABILITY OF DEATH FROM COVID (for under 60's)**

Serious Adverse Events

Deaths

A total of six (2 vaccine, 4 placebo) of 43,448 enrolled participants (0.01%) died during the reporting period from April 29, 2020 (first participant, first visit) to November 14, 2020 (cutoff date). Both vaccine recipients were >55 years of age; one experienced a cardiac arrest 62 days after vaccination #2 and died 3 days later, and the other died from arteriosclerosis 3 days after vaccination #1. The placebo recipients died from myocardial infarction (n=1), hemorrhagic stroke (n=1) or unknown causes (n=2); three of the four deaths occurred in the older group (>65 years of age). All deaths represent events that occur in the general population of the age groups where they occurred, at a similar rate.

Non-fatal SAEs

In the all-enrolled population of (total N=43,448), the proportions of participants who reported at least 1 SAE during the time period from Dose 1 to the data cutoff date (November 14, 2020) were 0.6% in the BNT162b2 vaccine group and 0.5% in the placebo group. The most common SAEs in the vaccine group which were numerically higher than in the placebo group were appendicitis (0.04%), acute myocardial infarction (0.02%), and cerebrovascular accident (0.02%), and in the placebo arm numerically higher than in the vaccine arm were pneumonia (0.03%), atrial fibrillation (0.02%), and syncope (0.02%). Occurrence of SAEs involving system organ classes and specific preferred terms were otherwise balanced between treatment groups, including no imbalance overall in cardiovascular serious adverse events.

Appendicitis was reported as a SAE for 12 participants, and numerically higher in the vaccine group: 8 vaccine participants (appendicitis [n=7], appendicitis perforated [n=1]) and 4 placebo participants (appendicitis [n=2], appendicitis perforated [n=1], complicated appendicitis [n=1]). All of the vaccine participants (n=8) and 2 placebo participants were younger than 65 years of age. The cases were considered unrelated to vaccination by the study investigators and occurred no more frequently than expected in the given age groups. FDA agrees that there is no clear basis upon which to suspect that this imbalance represents a vaccine-related risk.

According to Pfizer's own report 6 people in their study have died – shortly after receiving a vaccine. The total number of participants was 37796. 439 people were withdrawn from the study, for various reasons, 6 of which because they died.

So if we calculate the percentage of participants that died from the total number who participated: we get: $6 / 37363 * 100 = 0.016 \%$

This number is **higher** than the risk of death from covid to under 60 year olds based on the latest data from the NHS in the UK, which is **0.006 %**

If we calculate the number of people who had such serious adverse events that they had to be removed from the trial plus the number of deaths we get:

$$19 / 37376 * 100 = 0.05 \%$$

This number is almost 10 times higher than the risk of death from covid for an under 60 year old.

FEVER FEVER EVERYWHERE...

We recently reported that [Pfizer's Phase 2 trial had seen very high percentages of participants with dangerously high fevers](#). The latest reported Phase 3 figures are even more alarming:

Adverse Event	BNT162b2 Dose 1 N=2238 n (%)	Placebo Dose 1 N=2248 n (%)	BNT162b2 Dose 2 N=2045 n (%)	Placebo Dose 2 N=2053 n (%)
	Fever			
≥38.0°C	85 (3.7)	20 (0.9)	331 (15.8)	10 (0.5)
>38.0°C to 38.4°C	64 (2.8)	10 (0.4)	194 (9.2)	5 (0.2)
>38.4°C to 38.9°C	15 (0.7)	5 (0.2)	110 (5.2)	3 (0.1)
>38.9°C to 40.0°C	8 (0.3)	3 (0.1)	26 (1.2)	2 (0.1)
>40.0°C	0 (0.0)	2 (0.1)	1 (0.0)	0 (0.0)

Over 15% of participants in the 18-55 age group got a fever from the shot and **1.2 % HAD A DANGEROUS FEVER OF OVER 38.9°C TO 40°C.**

Let's remember that a fever of 40°C is potentially life threatening event.

Compare this again to the chance of dying from 'covid flu' which is 0.006 % :

According to Pfizer's own data – you will be 200 times MORE LIKELY to have a life endangering fever (over 38.9 °C to 40°C) than you would of ever dying from the covid flu.

You are also 10 times more likely to have either a severe adverse event or die from a Pfizer vaccine than you are to die from 'covid'.

You are 100 times more likely to have a Serious Adverse Reaction to Pfizer's vaccine than you are to die from covid if you are under 60.

Thanks for the report Pfizer, we are still waiting for your statement on why you need legal indemnity from being sued. We can only imagine.

Share this:



★ Like
One blogger likes this.

William J. Banks • December 9, 2020 • Covid, Science, World News

Published by William J. Banks

I have a University Certificate in Natural Sciences, (Biology Chemistry & Physics) and a University Diploma in Computer Science. [View more posts](#)

PREVIOUS POST

AP News Fails to Debunk Bill Gates' words: "there will be some risk and indemnification needed"

NEXT POST

Covid Burkas and Human Evolution

3 thoughts on "BOMBSHELL: More Death and Fever from Pfizer Vaccine than Covid"

Pingback: [BOMBSHELL: Pfizer Placebo blijkt geen placebo te zijn – 4 mensen overleden en een placebo? Axis World News – dezeweek.eu en Brandhaard.com](#)

Aschwin Wesselius
DECEMBER 10, 2020 AT 3:03 PM

Hi, interesting find! However, I can't spot in the PDF where it specifies the placebo as being actually a MenACWY vaccine. Can you disclose where you verified this to be true?
★ Like

Reply

William J. Banks
DECEMBER 10, 2020 AT 5:22 PM

Thanks Aschwin, yes, it clearly states it in the Phase 2 Lancet Paper:
[https://www.thelancet.com/journals/lancet/article/PIIS0140-6736\(20\)32466-1/fulltext](https://www.thelancet.com/journals/lancet/article/PIIS0140-6736(20)32466-1/fulltext)
"MenACWY was used as a comparator vaccine rather than a saline placebo to maintain masking of participants who had local or systemic reactions."
★ Like

Reply

Leave a Reply

Enter your comment here...