

COVID-19



- Get the latest public health information from CDC
- Get the latest research information from NIH | Español
- NIH staff guidance on coronavirus (NIH Only)

NIH RESEARCH MATTERS

January 26, 2021

Lasting immunity found after recovery from COVID-19


At a Glance

- The immune systems of more than 95% of people who recovered from COVID-19 had durable memories of the virus up to eight months after infection.
- The results provide hope that people receiving SARS-CoV-2 vaccines will develop similar lasting immune memories after vaccination.

After people recover from infection with a virus, the immune system retains a memory of it. Immune cells and proteins that circulate in the body can recognize and kill the pathogen if it's encountered again, protecting against disease and reducing illness severity.

This long-term immune protection involves several components. Antibodies—proteins that circulate in the blood—recognize foreign substances like viruses and neutralize them. Different types of T cells help recognize and kill pathogens. B cells make new antibodies when the body needs them.

All of these immune-system components have been found in people who recover from SARS-CoV-2, the virus that causes COVID-19. But the details of this immune response and how long it lasts after infection have been unclear. Scattered reports of reinfection with SARS-CoV-2 have raised concerns that the immune response to the virus might not be durable.



Colorized scanning electron micrograph of a cell, isolated from a patient sample, that is heavily infected with SARS-CoV-2 virus particles (red). *NIAID Integrated Research Facility, Fort Detrick, Maryland*

In this Edition

[Physician-pharmacist collaboration may improve care for opioid addiction](#)

[Acute heart transplant rejection detected earlier with new test](#)

[Lasting immunity found after recovery from COVID-19](#)

Search NIH Research Matters

Connect with Us

Subscribe to get NIH Research Matters by email

- [RSS Feed](#)
- [Facebook](#)
- [Email us](#)

Mailing Address:

To better understand immune memory of SARS-CoV-2, researchers led by Drs. Daniela Weiskopf, Alessandro Sette, and Shane Crotty from the La Jolla Institute for Immunology analyzed immune cells and antibodies from almost 200 people who had been exposed to SARS-CoV-2 and recovered.

Time since infection ranged from six days after symptom onset to eight months later. More than 40 participants had been recovered for more than six months before the study began. About 50 people provided blood samples at more than one time after infection.

The research was funded in part by NIH's National Institute of Allergy and Infectious Diseases (NIAID) and National Cancer Institute (NCI). Results were published on January 6, 2021, in *Science*.

The researchers found durable immune responses in the majority of people studied. Antibodies against the spike protein of SARS-CoV-2, which the virus uses to get inside cells, were found in 98% of participants one month after symptom onset. As seen in previous studies, the number of antibodies ranged widely between individuals. But, promisingly, their levels remained fairly stable over time, declining only modestly at 6 to 8 months after infection.

Virus-specific B cells increased over time. People had more memory B cells six months after symptom onset than at one month afterwards. Although the number of these cells appeared to reach a plateau after a few months, levels didn't decline over the period studied.

Levels of T cells for the virus also remained high after infection. Six months after symptom onset, 92% of participants had CD4+ T cells that recognized the virus. These cells help coordinate the immune response. About half the participants had CD8+ T cells, which kill cells that are infected by the virus.

As with antibodies, the numbers of different immune cell types varied substantially between individuals. Neither gender nor differences in disease severity could account for this variability. However, 95% of the people had at least 3 out of 5 immune-system components that could recognize SARS-CoV-2 up to 8 months after infection.

"Several months ago, our studies showed that natural infection induced a strong response, and this study now shows that the responses last," Weiskopf says. "We are hopeful that a similar pattern of responses lasting over time will also emerge for the vaccine-induced responses."

—by Sharon Reynolds

Related Links

- [Experimental Coronavirus Vaccine Highly Effective](https://www.nih.gov/news-events/nih-research-matters/experimental-coronavirus-vaccine-highly-effective) (https://www.nih.gov/news-events/nih-research-matters/experimental-coronavirus-vaccine-highly-effective)
- [Antibodies and T Cells Protect Against SARS-CoV-2](https://www.nih.gov/news-events/nih-research-matters/antibodies-t-cells-protect-against-sars-cov-2) (https://www.nih.gov/news-events/nih-research-matters/antibodies-t-cells-protect-against-sars-cov-2)

NIH Research Matters
Bldg. 31, Rm. 5B52,
MSC 2094
Bethesda, MD 20892-
2094

Popular Stories

Lasting immunity
found after recovery
from COVID-19

[Immune response to
vaccination after
COVID-19](#)

[Single-shot COVID-19
vaccine produces
immune response
against variants](#)

[Rehab program helps
frail older adults with
heart failure](#)

[Testing a blood
biomarker for
neurodegeneration](#)

About NIH Research Matters

Editor: Harrison Wein,
Ph.D. **Assistant**

Editor: Erin Bryant

**NIH Research
Matters** is a weekly
update of NIH
research highlights
reviewed by NIH's
experts. It's published
by the [Office of
Communications and
Public Liaison](#) in the
NIH Office of the
Director.

- [Immune Cells for Common Cold May Recognize SARS-CoV-2](https://www.nih.gov/news-events/nih-research-matters/immune-cells-common-cold-may-recognize-sars-cov-2) (https://www.nih.gov/news-events/nih-research-matters/immune-cells-common-cold-may-recognize-sars-cov-2)
- [Potent Neutralizing Antibodies Target New Regions of Coronavirus Spike](https://www.nih.gov/news-events/nih-research-matters/potent-neutralizing-antibodies-target-new-regions-coronavirus-spike) (https://www.nih.gov/news-events/nih-research-matters/potent-neutralizing-antibodies-target-new-regions-coronavirus-spike)
- [Potent Antibodies Found in People Recovered from COVID-19](https://www.nih.gov/news-events/nih-research-matters/potent-antibodies-found-people-recovered-covid-19) (https://www.nih.gov/news-events/nih-research-matters/potent-antibodies-found-people-recovered-covid-19)
- [Novel Coronavirus Structure Reveals Targets for Vaccines and Treatments](https://www.nih.gov/news-events/nih-research-matters/novel-coronavirus-structure-reveals-targets-vaccines-treatments) (https://www.nih.gov/news-events/nih-research-matters/novel-coronavirus-structure-reveals-targets-vaccines-treatments)
- [Coronavirus \(COVID-19\)](https://covid19.nih.gov/) (https://covid19.nih.gov/)
- [Coronavirus Prevention Network](https://www.coronaviruspreventionnetwork.org/) (https://www.coronaviruspreventionnetwork.org/)
- [Coronavirus \(COVID-19\)](https://www.coronavirus.gov/) (https://www.coronavirus.gov/)

References: [Immunological memory to SARS-CoV-2 assessed for up to 8 months after infection](#). Dan JM, Mateus J, Kato Y, Hastie KM, Yu ED, Faliti CE, Grifoni A, Ramirez SI, Haupt S, Frazier A, Nakao C, Rayaprolu V, Rawlings SA, Peters B, Krammer F, Simon V, Saphire EO, Smith DM, Weiskopf D, Sette A, Crotty S. *Science*. 2021 Jan 6:eabf4063. doi: 10.1126/science.abf4063. Online ahead of print. PMID: 33408181.

Funding: NIH's National Institute of Allergy and Infectious Diseases (NIAID) and National Cancer Institute (NCI); La Jolla Institute for Immunology; John and Mary Tu Foundation; Bill and Melinda Gates Foundation; Mastercard; Wellcome; Emergent Ventures; Collaborative Influenza Vaccine Innovation Centers; JPB Foundation; Cohen Foundation; Open Philanthropy Project.

NIH...Turning Discovery Into Health[®]

National Institutes of Health, 9000 Rockville Pike, Bethesda, Maryland 20892

[U.S. Department of Health and Human Services](#)